of the head ty of the RF mant at 0.8 spot'' in its dampen its nd yield a by us. uced by RF

ined by the irces of senly is needed erencing of

additional 1. The therubtless cors of irradiacularities of hold levels.

r colleagues, took part in

aracteristics Radio Sci.,

t of the 7th al Engineerolm.

ive hearing: a by pulsed

modulated 42.

se to modu-, 689-692. irradiation,

), A coaxial ng illumina-

rception of tromagnetic

nsen (1975), an auditory 4*cad. Sci.*, Lebovitz, R. M., and R. L. Seaman (1977), Microwave hearing: The response of single auditory neurons in the cat to pulsed microwave radiation, *Radio Sci.*, 12(6S), 229-236.

Lin, J. C. (1976), Theoretical analysis of microwave-generated auditory effects in animals and man, in Biological Effects of Electromagnetic Waves, Selected Papers of the USNC/URSI Annual Meeting, Boulder, Colorado, October 20-23, 1975, Vol. I, edited by C. C. Johnson and M. L. Shore, HEW Publ. FDA (77-8010), 36-48, U. S. Government Printing Office, Washington, D. C. 20402.

Tyazhelov, V. V., R. E. Tigranian, and E. P. Khizhniak (1977), New artifact-free electrodes for recording of biological potentials in strong electromagnetic fields, *Radio Sci.*, 12(6S), 121-124.